



Acetaminophen (Tylenol™, Feverall™) Dosage for Fever and Pain

- Review the medication concentration and select the column that matches from the chart below.
- Use **only** the enclosed dropper or medication cup that comes with the product. (Kitchen teaspoons are not accurate measures for medication.)
- Dose may be given every 4 hours. Do not use more than 5 times in 24 hours.

The following abbreviations are used on this dosage chart:

Milligram (mg)

Milliliter (mL)

Not applicable (--) : This form of medication should not be given to a child of this weight.

Acetaminophen (Tylenol™, Feverall™) Dosing Guideline					
Child's weight (pounds approx)	Liquid 160 mg/5 mL	Children's chewable or meltaway tablet 80 mg	Junior strength chewable or melt- away tablet 160 mg	Adult tablet 325 mg	Adult extra strength tablet 500 mg
2.7 to 5.4 kg (6 to 11 pounds)	1.25 mL (40 mg)	--	--	--	--
5.5 to 8.1 kg (12 to 17 pounds)	2.5 mL (80 mg)	1 tablet (80 mg)	--	--	--
8.2 to 10.8 kg (18 to 23 pounds)	3.75 mL (120 mg)	1 ½ tablets (120 mg)	--	--	--
10.9 to 16.3 kg (24 to 35 pounds)	5 mL (160 mg)	2 tablets (160 mg)	1 tablet (160 mg)	--	--
16.4 to 21.7 kg (36 to 47 pounds)	7.5 mL (240 mg)	3 tablets (240 mg)	1 ½ tablets (240 mg)	--	--
21.8 to 27.2 kg (48 to 59 pounds)	10 mL (320 mg)	4 tablets (320 mg)	2 tablets (320 mg)	1 tablet (325 mg)	--
27.3 to 32.6 kg (60 to 71 pounds)	12.5 mL (400 mg)	5 tablets (400 mg)	2 ½ tablets (400 mg)	1 tablet (325 mg)	--
32.7 to 43.1 kg (72 to 95 pounds)	15 mL (480 mg)	6 tablets (480 mg)	3 tablets (480 mg)	1 ½ tablets (487.5 mg)	1 tablet (500 mg)
43.2 to 66.4 kg (95 to 146 pounds)	20 mL (640 mg)	8 tablets (640 mg)	4 tablets (640 mg)	2 tablets (650 mg)	1 tablet (500 mg)