



# Pediatric Health Promotion Visit and Vaccination Schedule

Each year	Influenza.....	Vaccination only
All ≥6 months	COVID-19 vaccination series.....	Vaccination only
Newborn	HepB.....	At the hospital
3–5 days	.....	Encounter
2 weeks–1 month	.....	Encounter Optional
2 months	PCV13, RV5, DTaP-IPV-Hib-Hepb.....	Encounter
4 months	PCV13, RV5, DTaP-IPV-Hib-Hepb.....	Encounter
6 months	PCV13, RV5, DTaP-IPV-Hib-Hepb.....	Encounter/Alternative Option
9 months	.....	Encounter
12 months	HepA, MMR, VAR.....	Encounter/Alternative Option
15 months	PCV13, DTaP-IPV/Hib.....	Encounter
18 months	.....	Encounter
2 years	HepA.....	Encounter
30 months	.....	Encounter
3 years	.....	Encounter
4 years	.....	Encounter
5 years	DTaP-IPV, MMRV.....	Encounter
6 years	.....	Encounter
7 years	.....	Encounter/Alternative Option
8 years	.....	Encounter
9 years	HPV (2 doses 6 months apart).....	Encounter/Alternative Option
10 years	.....	Encounter
11 years	MCV4, Tdap.....	Encounter
12 years	.....	Encounter
13 years	.....	Encounter
14 years	.....	Encounter
15 years	.....	Encounter
16 years	MCV4, MenB* (optional 2 doses 1 month apart).....	Encounter
17 years	.....	Encounter
18 years	.....	Encounter
19 years	.....	Encounter
20 years	.....	Encounter
21 years	.....	Encounter

Visit Definitions and Full Vaccine Names on back.

## *Pediatric Health Promotion Visit and Vaccination Schedule* (continued)

### **Visit Definitions**

Encounter	Well child visit face-to-face with a health care provider and care team.
Encounter Optional	Provider, care team member, and parent decide together at time of 3 to 5 days well visit whether the newborn needs a visit at 2 or 4 weeks of age.
Alternative Option	Health promotion visit can be a face-to-face encounter with a health care team member (for example, a nurse well visit).

### **Full Vaccine Names**

DTaP	Diphtheria-Tetanus-acellular Pertussis vaccine (pediatric form)
DTaP-IPV	Diphtheria-Tetanus-acellular Pertussis vaccine-Inactivated Poliovirus Vaccine (Quadracel)
DTaP-IPV/Hib	Diphtheria-Tetanus-acellular Pertussis vaccine-Inactivated Poliovirus Vaccine/Haemophilus influenzae type B vaccine (Pentacel)
DTaP-IPV-Hib-Hepb	Diphtheria-Tetanus-acellular Pertussis-Inactivated Poliovirus-Haemophilus influenzae type b-Hepatitis B vaccine (Vaxelis)
HepA	Hepatitis A vaccine (minimum spacing 6 months between doses)
HepB	Hepatitis B vaccine
HPV	Human Papillomavirus Vaccine (2 doses 6 months apart for immunocompetent patients who get the first dose before 15 years of age; otherwise 3 doses at 0, 2, and 6 months)
IPV	Inactivated Poliovirus Vaccine
MCV4	Meningococcal Conjugate Vaccine (quadrivalent)
*MenB	Meningococcal B vaccine (optional) 16 through 23 years of age; (Bexsero – 2 doses at least 1 month apart)
MMR	Measles-Mumps-Rubella vaccine
MMRV	Measles-Mumps-Rubella-Varicella vaccine (give MMR and VAR separately before age 4 or after age 12 years)
PCV13	Pneumococcal Conjugate Vaccine (13-valent)
RV5	Rotavirus (pentavalent) vaccine (first dose no later than 14 weeks, 6 days of age and last dose no later than 8 months, 0 days of age)
Tdap	Tetanus-diphtheria-acellular pertussis vaccine (adolescent-adult form)
VAR	Varicella zoster vaccine (chicken pox)